

DAUGAVET CENTERS AND DIRECT SUMS OF BANACH SPACES

TETIANA V. BOSENKO

ABSTRACT. A linear continuous nonzero operator $G: X \rightarrow Y$ is a Daugavet center if every rank-1 operator $T: X \rightarrow Y$ satisfies $\|G + T\| = \|G\| + \|T\|$. We study the case when either X or Y is a sum $X_1 \oplus_F X_2$ of two Banach spaces X_1 and X_2 by some two-dimensional Banach space F . We completely describe the class of those F such that for some spaces X_1 and X_2 there exists a Daugavet center acting from $X_1 \oplus_F X_2$, and the class of those F such that for some pair of spaces X_1 and X_2 there is a Daugavet center acting into $X_1 \oplus_F X_2$. We also present several examples of such Daugavet centers.

1. INTRODUCTION

In the present paper we consider real Banach spaces which do not equal $\{0\}$, and denote them E , X or Y . A linear continuous nonzero operator $G: X \rightarrow Y$ is called a *Daugavet center* [3] if every rank-1 operator $T: X \rightarrow Y$ satisfies the equation

$$(1.1) \quad \|G + T\| = \|G\| + \|T\|.$$

Definition 1.1. We say that X is a *Daugavet domain* if there exists a Daugavet center $G: X \rightarrow Y$ for some Y , and is a *Daugavet range* if there is a Daugavet center $G: E \rightarrow X$ for some E .

Throughout this paper $F = (\mathbb{R}^2, \|\cdot\|)$ with $\|(1, 0)\| = \|(0, 1)\| = 1$ and

$$(1.2) \quad \|(a_1, a_2)\| = \|(|a_1|, |a_2|)\|$$

for every $(a_1, a_2) \in F$. For Banach spaces X_1 and X_2 their F -sum $X_1 \oplus_F X_2$ is the space of all pairs (x_1, x_2) where $x_1 \in X_1$ and $x_2 \in X_2$, $\|(x_1, x_2)\| := \|(\|x_1\|, \|x_2\|)\|_F$.

We introduce the following order on F : $(a_1, a_2) \geq (b_1, b_2)$ if $a_1 \geq b_1$ and $a_2 \geq b_2$. It follows from (1.2) and a convexity argument that for every $(a_1, a_2), (b_1, b_2) \in F$ with $(|a_1|, |a_2|) \leq (|b_1|, |b_2|)$ the inequality $\|(a_1, a_2)\| \leq \|(b_1, b_2)\|$ holds true. In this partial order F is a Banach lattice [8], so we will use the term “two-dimensional lattice” for F in the sequel.

The problem which we solve in this paper, consists of two parts: first, we characterize the class of those F for which there exist X_1 and X_2 such that

2000 *Mathematics Subject Classification.* Primary 46B04; secondary 46B20, 46B40.

Key words and phrases. Daugavet center, Daugavet property, direct sum of Banach spaces.

$X_1 \oplus_F X_2$ is a Daugavet domain, and secondly, we characterize the class of those F for which there are X_1 and X_2 such that $X_1 \oplus_F X_2$ is a Daugavet range.

Remark that a Daugavet domain and a Daugavet range are generalizations of a Banach space with the Daugavet property, and this motivates our interest in the subject. A Banach space X is said to have the Daugavet property if the identity operator $\text{Id}: X \rightarrow X$ is a Daugavet center. The study of spaces with the Daugavet property is a rapidly developing branch of Banach space theory (see [6], [11], [13], and the most recent developments in [5], [7]). The following classical spaces have the Daugavet property: $C(K)$ where K is a compact without isolated points [4], $L_1(\mu)$ and $L_\infty(\mu)$ where μ has no atoms [9], and many Banach algebras ([14], [15]). Some exotic spaces have the Daugavet property as well, for instance, Talagrand's space ([6], [12]) and Bourgain-Rosenthal's space ([2], [7]).

Let us recall some recent results [3] related to Daugavet centers. If G is a Daugavet center then (1.1) also holds true when T is a strong Radon-Nikodým operator, e.g., a weakly compact operator. If X is a Daugavet domain or a Daugavet range then X contains subspaces isomorphic to ℓ_1 , is non-reflexive and does not have an unconditional basis (countable or uncountable). One cannot even embed such an X into a space having an unconditional basis or having a representation as unconditional sum of reflexive subspaces. In [10] Popov proves that every isometric embedding of $L_1[0, 1]$ into itself is a Daugavet center. However, in [3] one can find examples of Daugavet centers which are not isometries.

The present work is inspired by [1]. It was shown in [1] and [6] that if X_1 and X_2 have the Daugavet property and $F = \ell_1^{(2)}$ or $\ell_\infty^{(2)}$ then $X_1 \oplus_F X_2$ has the Daugavet property as well. In [1] the authors prove that $X_1 \oplus_F X_2$ has the Daugavet property only if $F = \ell_1^{(2)}$ or $\ell_\infty^{(2)}$. In our paper we generalize these results of [1], but use a new approach to the problem. Surprisingly in both parts of our problem we discover other spaces apart from $F = \ell_1^{(2)}$ and $F = \ell_\infty^{(2)}$, which satisfy our demands.

Our approach is based on a necessary condition for a general Banach space X to be a Daugavet domain and on a necessary condition for X to be a Daugavet range. We deduce these two conditions in Section 2 (see Definition 2.1, Lemma 2.5 and Definition 2.2, Lemma 2.6) and then we show in Section 3 how they depend on F when $X = X_1 \oplus_F X_2$ (see Lemma 3.6 and Lemma 3.7).

In Section 4 we find a rather small class \mathfrak{N}_2 such that if $X_1 \oplus_F X_2$ is a Daugavet range then $F \in \mathfrak{N}_2$, and in Section 5 we discover the analogous class \mathfrak{M}_2 for the case of a Daugavet domain. Then for every $F \in \mathfrak{M}_2$ we present an example of a Daugavet center acting *from* a sum of two Banach spaces by F (see Proposition 6.2), and this solves the first part of our problem. In a very similar way we solve its second part, namely we give examples of Daugavet centers acting *into* a sum of two Banach spaces by F

for every $F \in \mathfrak{N}_2$ (see Proposition 6.6). The obtained results illustrate that the notions of a Daugavet domain and a Daugavet range do not refer to the same Banach spaces.

Throughout this paper B_X denotes the closed unit ball of X and S_X denotes its unit sphere. We use the notation

$$B_F^+ := \{a \in B_F : a \geq 0\}$$

for the positive part of the unit ball and

$$S_F^+ := \{a \in S_F : a \geq 0\}$$

for the positive part of the unit sphere of F . We denote

$$S(B_X, z^*, \varepsilon) := \{x \in B_X : z^*(x) > 1 - \varepsilon\}$$

the slice of B_X determined by $z^* \in S_{X^*}$ and $\varepsilon > 0$.

$$S(B_{X^*}, z, \varepsilon) = \{x^* \in B_{X^*} : x^*(z) > 1 - \varepsilon\}$$

denotes the weak* slice of B_{X^*} determined by $z \in S_X$ and $\varepsilon > 0$. For an $x^* \in X^*$ and a $y \in Y$ the symbol $x^* \otimes y$ stands for the operator which acts from X into Y as follows: $(x^* \otimes y)(x) = x^*(x)y$.

Finally, let us cite a fact that we frequently use in the sequel.

Theorem 1.2 ([3], Theorem 2.1). *For an operator $G: X \rightarrow Y$ with $\|G\| = 1$ the following assertions are equivalent:*

- (i) G is a Daugavet center.
- (ii) For every $y_0 \in S_Y$, $x_0^* \in S_{X^*}$ and $\varepsilon > 0$ there is an $x \in S(B_X, x_0^*, \varepsilon)$ with $\|Gx + y_0\| > 2 - \varepsilon$.
- (iii) For every $y_0 \in S_Y$, $x_0^* \in S_{X^*}$ and $\varepsilon > 0$ there is a $y^* \in S(B_{Y^*}, y_0, \varepsilon)$ with $\|G^*y^* + x_0^*\| > 2 - \varepsilon$.

2. BANACH SPACES DENYING THE DAUGAVET PROPERTY

Definition 2.1. *We say that X denies the Daugavet property with a set $A \subset S_X$ if there is an $\varepsilon > 0$ such that for every $y \in A$ there exists an $x^* \in S_{X^*}$ satisfying*

$$(2.1) \quad \|\text{Id} + x^* \otimes y\| < 2 - \varepsilon.$$

Definition 2.2. *We say that X star-denies the Daugavet property with a set $A \subset S_{X^*}$ if there is an $\varepsilon > 0$ such that for every $x^* \in A$ there exists a $y \in S_X$ satisfying (2.1).*

Lemma 2.3. *For $A \subset S_X$ the following assertions are equivalent:*

- (i) X denies the Daugavet property with A .
- (ii) There is an $\varepsilon > 0$ such that for every $y \in A$ a functional $x^* \in S_{X^*}$ may be chosen so that every $x \in S(B_X, x^*, \varepsilon)$ fulfills $\|x + y\| < 2 - \varepsilon$.

- (iii) *There is an $\varepsilon > 0$ such that for every $y \in A$ a functional $x^* \in S_{X^*}$ may be chosen so that every $y^* \in S(B_{X^*}, y, \varepsilon)$ fulfills $\|x^* + y^*\| < 2 - \varepsilon$.*

Proof. (i) \Rightarrow (ii) We have that there is an $\varepsilon > 0$ such that for every $y \in A$ there exists an $x^* \in S_{X^*}$ satisfying

$$\|\text{Id} + x^* \otimes y\| = \sup_{x \in B_X} \|x + x^*(x)y\| < 2 - \varepsilon.$$

Hence $\|x + x^*(x)y\| < 2 - \varepsilon$ for every $x \in B_X$. Let $x \in S(B_X, x^*, \varepsilon/2)$ then

$$\|x + y\| \leq \|x + x^*(x)y\| + \|y - x^*(x)y\| < 2 - \varepsilon + |1 - x^*(x)| \cdot \|y\| < 2 - \varepsilon + \frac{\varepsilon}{2} = 2 - \frac{\varepsilon}{2}$$

which implies (ii).

(ii) \Rightarrow (i) Let ε and x^* be from (ii). It is sufficient to show that $\|x + x^*(x)y\| \leq 2 - \varepsilon/2$ for every $x \in B_X$. Let $x \in S(B_X, x^*, \varepsilon/2)$ then

$$\|x + x^*(x)y\| \leq \|x + y\| + \|y - x^*(x)y\| < 2 - \varepsilon + |1 - x^*(x)| \cdot \|y\| < 2 - \frac{\varepsilon}{2}.$$

Let $x \in S(B_X, -x^*, \varepsilon/2)$ then $-x \in S(B_X, x^*, \varepsilon/2)$. Hence $\|x - y\| < 2 - \varepsilon$ and

$$\|x + x^*(x)y\| \leq \|x - y\| + \|y + x^*(x)y\| < 2 - \varepsilon + |1 + x^*(x)| \cdot \|y\| < 2 - \frac{\varepsilon}{2}.$$

Finally, let $x \in B_X \setminus (S(B_X, x^*, \varepsilon/2) \cup S(B_X, -x^*, \varepsilon/2))$ then

$$\|x + x^*(x)y\| \leq \|x\| + |x^*(x)| \cdot \|y\| \leq 2 - \frac{\varepsilon}{2}.$$

The equivalence (i) \Leftrightarrow (iii) can be proved in a very similar fashion to (i) \Leftrightarrow (ii) using the fact that the norms of an operator and of its adjoint coincide. \square

Lemma 2.4. *For $A \subset S_{X^*}$ the following assertions are equivalent:*

- (i) *X star-denies the Daugavet property with A .*
- (ii) *There is an $\varepsilon > 0$ such that for every $x^* \in A$ a vector $y \in S_X$ may be chosen so that every $x \in S(B_X, x^*, \varepsilon)$ fulfills $\|x + y\| < 2 - \varepsilon$.*
- (iii) *There is an $\varepsilon > 0$ such that for every $x^* \in A$ a vector $y \in S_X$ may be chosen so that every $y^* \in S(B_{X^*}, y, \varepsilon)$ fulfills $\|x^* + y^*\| < 2 - \varepsilon$.*

One can prove Lemma 2.4 the same way as Lemma 2.3. The following two lemmas form the main result of this section.

Lemma 2.5. *Let there exist $\delta > 0$ and $z^* \in S_{X^*}$ such that X denies the Daugavet property with $S(B_X, z^*, \delta) \cap S_X$. Then X is not a Daugavet domain.*

Proof. According to Definition 1.1 we must prove that any $G: X \rightarrow Y$ is not a Daugavet center for any Y . It is easy to see that if G is a Daugavet center then $G/\|G\|$ is as well, so we consider only the case $\|G\| = 1$.

Take the ε from item (ii) of Lemma 2.3. At first we show that if every $z \in S(B_X, z^*, \delta)$ satisfies $\|Gz\| \leq 1 - \varepsilon/2$ then G is not a Daugavet center.

Put $\varepsilon_0 := \min\{\varepsilon/2, \delta\}$, then for every $y \in S_Y$ and every $z \in S(B_X, z^*, \varepsilon_0)$ we have

$$\|y + Gz\| \leq 1 + \|Gz\| \leq 2 - \frac{\varepsilon}{2} \leq 2 - \varepsilon_0.$$

Theorem 1.2, item (ii) implies that G is not a Daugavet center.

So, we suppose that there is a $z_0 \in S(B_X, z^*, \delta)$ with

$$(2.2) \quad \|Gz_0\| > 1 - \frac{\varepsilon}{2}.$$

We can assume $\|z_0\| = 1$, because if $z_0 \in B_X$ fulfills $z^*(z_0) > 1 - \delta$ and (2.2) then $z_0/\|z_0\|$ does as well. In addition, (2.2) implies that there is a $y_0 \in S_Y$ with $\|y_0 - Gz_0\| < \varepsilon/2$. Since X denies the Daugavet property with $S(B_X, z^*, \delta) \cap S_X$, there is an $x^* \in S_{X^*}$ such that every $x \in S(B_X, x^*, \varepsilon)$ satisfies $\|x + z_0\| < 2 - \varepsilon$. Hence for every $x \in S(B_X, x^*, \varepsilon)$ we have

$$\|y_0 + Gx\| \leq \|y_0 - Gz_0\| + \|Gx + Gz_0\| < \frac{\varepsilon}{2} + \|x + z_0\| < 2 - \frac{\varepsilon}{2}.$$

By Theorem 1.2, item (ii) G is not a Daugavet center. \square

Lemma 2.6. *Let there exist $\delta > 0$ and $z \in S_X$ such that X star-denies the Daugavet property with $S(B_{X^*}, z, \delta) \cap S_{X^*}$. Then X is not a Daugavet range.*

Using item (iii) of Lemma 2.4 and item (iii) of Theorem 1.2 one can prove Lemma 2.6 in a very similar fashion to Lemma 2.5.

3. TWO-DIMENSIONAL LATTICES DENYING THE POSITIVE DAUGAVET PROPERTY

Definition 3.1. *We say that F denies the positive Daugavet property with $A \subset S_F^+$ if there is an $\varepsilon > 0$ such that for every $a \in A$ there exists an $f^* \in S_{F^*}^+$ satisfying*

$$(3.1) \quad \|\text{Id} + f^* \otimes a\| < 2 - \varepsilon.$$

Definition 3.2. *We say that F star-denies the positive Daugavet property with $A \subset S_{F^*}^+$ if there is an $\varepsilon > 0$ such that for every $f^* \in A$ there exists an $a \in S_F^+$ satisfying (3.1).*

The following two lemmas are complete analogs of Lemmas 2.3 and 2.4, so we skip their proofs.

Lemma 3.3. *For $A \subset S_F^+$ the following assertions are equivalent:*

- (i) *F denies the positive Daugavet property with A .*
- (ii) *There is an $\varepsilon > 0$ such that for every $a \in A$ a functional $f^* \in S_{F^*}^+$ may be chosen so that every $b \in S(B_F, f^*, \varepsilon) \cap B_F^+$ fulfills $\|a + b\| < 2 - \varepsilon$.*
- (iii) *There is an $\varepsilon > 0$ such that for every $a \in A$ a functional $f^* \in S_{F^*}^+$ may be chosen so that every $g^* \in S(B_{F^*}, a, \varepsilon) \cap B_{F^*}^+$ fulfills $\|f^* + g^*\| < 2 - \varepsilon$.*

Lemma 3.4. *For $A \subset S_{F^*}^+$ the following assertions are equivalent:*

- (i) *F star-denies the positive Daugavet property with A .*
- (ii) *There is an $\varepsilon > 0$ such that for every $f^* \in A$ a vector $a \in S_F^+$ may be chosen so that every $b \in S(B_F, f^*, \varepsilon) \cap B_F^+$ fulfills $\|a + b\| < 2 - \varepsilon$.*
- (iii) *There is an $\varepsilon > 0$ such that for every $f^* \in A$ a vector $a \in S_F^+$ may be chosen so that every $g^* \in S(B_{F^*}, a, \varepsilon) \cap B_{F^*}^+$ fulfills $\|f^* + g^*\| < 2 - \varepsilon$.*

Recall that $F^* = \mathbb{R}^2$ with the norm

$$\|(f_1, f_2)\|_{F^*} := \max_{(a_1, a_2) \in B_F} |f_1 a_1 + f_2 a_2|$$

and $F^{**} = F$. We introduce an order on F^* the same way as on F . It is easy to see that $\|(1, 0)\|_{F^*} = \|(0, 1)\|_{F^*} = 1$ and $\|(f_1, f_2)\|_{F^*} = \|(|f_1|, |f_2|)\|_{F^*}$ for every $(f_1, f_2) \in F^*$. Hence F^* is a two-dimensional lattice as well. Lemmas 3.3 and 3.4 evidently imply the following fact (which one can easily deduce from Definitions 3.1 and 3.2 as well).

Lemma 3.5. *Let $A \subset S_F^+$ and $\tilde{A} \subset S_{F^*}^+$.*

- (a) *If F denies the positive Daugavet property with A then F^* star-denies the positive Daugavet property with A .*
- (b) *If F star-denies the positive Daugavet property with \tilde{A} then F^* denies the positive Daugavet property with \tilde{A} .*

Here is the key lemma of this section. In its proof we use the idea from Theorem 5.1 of [1].

Lemma 3.6. *Let there exist $w^* \in S_{F^*}^+$ and $\delta > 0$ such that F denies the positive Daugavet property with $S(B_F, w^*, \delta) \cap S_F^+$. Then $X_1 \oplus_F X_2$ is not a Daugavet domain for any X_1 and X_2 .*

Proof. It is easy to see that $(X_1 \oplus_F X_2)^* = X_1^* \oplus_{F^*} X_2^*$ for every X_1 and X_2 . Pick a $z^* = (z_1^*, z_2^*) \in S_{(X_1 \oplus_F X_2)^*}$ with $(\|z_1^*\|, \|z_2^*\|) = w^*$. Then for a $y = (y_1, y_2) \in S(B_{X_1 \oplus_F X_2}, z^*, \delta) \cap S_{X_1 \oplus_F X_2}$ we have

$$\|z_1^*\| \|y_1\| + \|z_2^*\| \|y_2\| \geq z_1^*(y_1) + z_2^*(y_2) = z^*(y) > 1 - \delta.$$

Hence $a := (\|y_1\|, \|y_2\|) \in S(B_F, w^*, \delta) \cap S_F^+$. By item (ii) of Lemma 3.3 there exist $\varepsilon > 0$ and $f^* \in S_{F^*}^+$ such that every $b \in S(B_F, f^*, \varepsilon) \cap B_F^+$ satisfies $\|a + b\| < 2 - \varepsilon$.

Pick an $x^* = (x_1^*, x_2^*) \in S_{(X_1 \oplus_F X_2)^*}$ with $(\|x_1^*\|, \|x_2^*\|) = f^*$. Then for every $x = (x_1, x_2) \in S(B_{X_1 \oplus_F X_2}, x^*, \varepsilon)$ we have $b_x := (\|x_1\|, \|x_2\|) \in S(B_F, f^*, \varepsilon) \cap B_F^+$ and therefore

$$\begin{aligned} \|x + y\| &= \|(\|x_1 + y_1\|, \|x_2 + y_2\|)\| \\ &\leq \|(\|x_1\| + \|y_1\|, \|x_2\| + \|y_2\|)\| = \|a + b_x\| < 2 - \varepsilon. \end{aligned}$$

By item (ii) of Lemma 2.3 $X_1 \oplus_F X_2$ denies the Daugavet property for $S(B_{X_1 \oplus_F X_2}, z^*, \delta) \cap S_{X_1 \oplus_F X_2}$. So, Lemma 2.5 implies that $X_1 \oplus_F X_2$ is not a Daugavet domain. \square

The same conclusions based on item (iii) of Lemma 3.4, item (iii) of Lemma 2.4, and Lemma 2.6 prove the following fact:

Lemma 3.7. *Let there exist $w \in S_F^+$ and $\delta > 0$ such that F star-denies the positive Daugavet property with $S(B_{F^*}, w, \delta) \cap S_{F^*}^+$. Then $X_1 \oplus_F X_2$ is not a Daugavet range for any X_1 and X_2 .*

4. SUMS OF SPACES WHICH ARE NOT DAUGAVET RANGES

In this section we find a large class of those F which star-deny the positive Daugavet property with some $S(B_{F^*}, w, \delta) \cap S_{F^*}^+$. Throughout this and the following sections $e_1 := (1, 0) \in S_F^+$, $e_2 := (0, 1) \in S_F^+$, and the symbol $[a, b]$ is reserved for the line segment with the end points in $a, b \in F$.

Lemma 4.1. *Let D be a closed subset of $S_{F^*}^+$. Suppose for every $f^* \in D$ there exists an $\varepsilon > 0$ such that the property $P(f^*, \varepsilon) := \{\text{there is an } a \in S_F^+ \text{ such that every } b \in S(B_F, f^*, \varepsilon) \cap B_F^+ \text{ satisfies } \|a + b\| < 2 - \varepsilon\}$ holds true. Then F star-denies the positive Daugavet property with D .*

Proof. Note that if $P(f^*, \varepsilon)$ holds true then $P(f^*, \varepsilon_1)$ holds for every $\varepsilon_1: 0 < \varepsilon_1 < \varepsilon$. Our goal is to show that there exists a common $\varepsilon_{\min} > 0$ such that $P(f^*, \varepsilon_{\min})$ holds true for every $f^* \in D$.

Consider the function $u(f^*): D \rightarrow (0, 1)$, $u(f^*) = \sup\{\varepsilon > 0: P(f^*, \varepsilon) \text{ holds true}\}$. Let us prove that $u(f^*)$ reaches its minimum value on D . Since D is compact, it is sufficient to show that $u(f^*)$ is lower semicontinuous, i.e. that the set $u^{-1}((x, 1))$ is open for every $x \in [0, 1)$.

Let $f^* \in u^{-1}((x, 1))$. This means that $u(f^*) = \sup\{\varepsilon > 0: P(f^*, \varepsilon) \text{ holds true}\} > x$. Hence there exist $\varepsilon_0 > x$ and $a \in S_F^+$ such that every $b \in S(B_F, f^*, \varepsilon_0) \cap B_F^+$ fulfills $\|a + b\| < 2 - \varepsilon_0$.

Take an $\varepsilon_1: x < \varepsilon_1 < \varepsilon_0$ and put $\delta := \varepsilon_0 - \varepsilon_1$. The set $D \cap B_{F^*}(f^*, \delta)$ is a relative neighborhood of f^* in D . Let us show that $D \cap B_{F^*}(f^*, \delta) \subset u^{-1}((x, 1))$.

Let $f_1^* \in D \cap B_{F^*}(f^*, \delta)$. Then every $b \in S(B_F, f_1^*, \varepsilon_1) \cap B_F^+$ fulfills

$$f^*(b) \geq f_1^*(b) - \delta > 1 - \varepsilon_1 - \delta = 1 - \varepsilon_0.$$

Thus $b \in S(B_F, f^*, \varepsilon_0) \cap B_F^+$, so we have

$$\|a + b\| < 2 - \varepsilon_0 < 2 - \varepsilon_1.$$

This means that $u(f_1^*) \geq \varepsilon_1 > x$ and $f_1^* \in u^{-1}((x, 1))$. Consequently, $u^{-1}((x, 1))$ is open and $u(f^*)$ is lower semicontinuous. Then there exists an $f_0^* \in D$ such that

$$u(f_0^*) = \min_{f^* \in D} u(f^*).$$

Take an $\varepsilon_{\min}: 0 < \varepsilon_{\min} < u(f_0^*)$ then $P(f^*, \varepsilon_{\min})$ holds true for every $f^* \in D$. \square

Lemma 4.2. *Let $a \in S_F^+$ and $f^* \in S_{F^*}^+$. Suppose for every $\varepsilon > 0$ there is a $b \in S(B_F, f^*, \varepsilon) \cap B_F^+$ with $\|a + b\| \geq 2 - \varepsilon$. Then there exists a $b_0 \in S_F^+$ such that $f^*(b_0) = 1$ and $[a, b_0] \subset S_F^+$.*

Proof. Consider a vanishing sequence $\{\varepsilon_n\}_{n=1}^\infty$, $\varepsilon_n > 0$. For every $n \in \mathbb{N}$ there exists a $b_n \in B_F^+$ with $f^*(b_n) > 1 - \varepsilon_n$ and $\|a + b_n\| \geq 2 - \varepsilon_n$.

Since B_F^+ is a compact set, there exists a subsequence $\{b_{n_i}\}_{i=1}^\infty$ of $\{b_n\}_{n=1}^\infty$ that converges to some $b_0 \in B_F^+$. Then $f^*(b_0) = 1$ and $\|a + b_0\| = 2$ which implies $[a, b_0] \subset S_F^+$. \square

Denote \mathfrak{N}_3 the class of those F whose S_F^+ is a polygon which consists of at most three edges.

Lemma 4.3. *Let $F \notin \mathfrak{N}_3$. Then F star-denies the positive Daugavet property with $S_{F^*}^+$.*

Proof. Assume to the contrary that there exists an $f^* \in S_{F^*}^+$ such that for every $\varepsilon > 0$ and $a_0 \in S_F^+$ there is a $b \in S(B_F, f^*, \varepsilon) \cap B_F^+$ with $\|a_0 + b\| \geq 2 - \varepsilon$.

Consider the set $\Delta := \{a \in S_F^+ : f^*(a) = 1\}$. It is easy to see that Δ is a segment or a point. Put $a_0 := e_1$. By Lemma 4.2 there exists a $b_0 \in \Delta$ such that $[b_0, e_1] \subset S_F^+$. If we put $a_0 := e_2$ we obtain a $b_1 \in \Delta$ with $[b_1, e_2] \subset S_F^+$. Then $F \in \mathfrak{N}_3$, because S_F^+ consists of at most three segments: $[b_0, e_1]$, Δ and $[b_1, e_2]$. This contradiction completes the proof. \square

Lemma 4.4. *Let S_F^+ be a polygon which consists of exactly three edges. Then there exists a $w^* = (w_1, w_2) \in S_{F^*}^+$ with $w_1 < 1$ and $w_2 < 1$ such that F star-denies the positive Daugavet property with $S_{F^*}^+ \setminus \overset{\circ}{B}_{F^*}(w^*, \delta_0)$ for every $\delta_0 > 0$.*

Proof. Since S_F^+ consists of three edges, it has four vertexes. The points e_1 and e_2 are two of them, denote h_1 and h_2 the remaining ones in such a way that $[e_1, h_1] \cup [e_2, h_2] \subset S_F^+$. There is the unique $w^* = (w_1, w_2) \in S_{F^*}^+$ such that $\{a \in S_F^+ : w^*(a) = 1\} = [h_1, h_2]$. It is obvious that $w_1 < 1$ and $w_2 < 1$.

Consider a $\delta_0 > 0$ and an $f^* \in S_{F^*}^+ \setminus \overset{\circ}{B}_{F^*}(w^*, \delta_0)$. Denote $\Delta := \{a \in S_F^+ : f^*(a) = 1\}$, it is a segment or a point. Assume that for every $\varepsilon > 0$ and $a \in S_F^+$ there exists a $b \in S(B_F, f^*, \varepsilon) \cap B_F^+$ with $\|a + b\| \geq 2 - \varepsilon$. By Lemma 4.2 there are $b_1, b_2 \in \Delta$ such that $[b_1, e_1] \subset S_F^+$ and $[b_2, e_2] \subset S_F^+$. Hence $b_1 \in [e_1, h_1]$ and $b_2 \in [e_2, h_2]$. Since $[e_1, h_1] \cap [e_2, h_2] = \emptyset$ then $\Delta \not\subset [e_1, h_1]$, $\Delta \not\subset [e_2, h_2]$, and Δ is a segment. Consequently, $\Delta \subset [h_1, h_2]$. But then $w^* = f^*$, so we come to contradiction.

Thus for every $f^* \in S_{F^*}^+ \setminus \overset{\circ}{B}_{F^*}(w^*, \delta_0)$ there are $\varepsilon > 0$ and $a \in S_F^+$ such that every $b \in S(B_F, f^*, \varepsilon) \cap B_F^+$ satisfies $\|a + b\| < 2 - \varepsilon$. Since $S_{F^*}^+ \setminus \overset{\circ}{B}_{F^*}(w^*, \delta_0)$ is closed, Lemma 4.1 implies the needed result. \square

Denote \mathfrak{N}_2 the class of those F whose S_F^+ is a polygon which consists of at most two edges.

Corollary 4.5. *Let $F \notin \mathfrak{N}_2$. Then there is a $\delta > 0$ such that F star-denies the positive Daugavet property with $S(B_{F^*}, e_1, \delta) \cap S_{F^*}^+$.*

Proof. If $F \notin \mathfrak{N}_3$ then by Lemma 4.3 the statement is proved.

If S_F^+ is a polygon which consists of exactly three edges then by Lemma 4.4 there exists a $w^* = (w_1, w_2) \in S_{F^*}^+$ with $w_1 < 1$ such that F star-denies the positive Daugavet property with $S_{F^*}^+ \setminus \mathring{B}_{F^*}(w^*, \delta_0)$ for every $\delta_0 > 0$. Pick a $\delta_0 > 0$ with $\delta_0 + w_1 < 1$ and a δ such that $0 < \delta < 1 - w_1 - \delta_0$. Then every $f^* \in \mathring{B}_{F^*}(w^*, \delta_0)$ satisfies

$$f^*(e_1) < w^*(e_1) + \delta_0 = w_1 + \delta_0 < 1 - \delta.$$

Hence $\mathring{B}_{F^*}(w^*, \delta_0) \cap S(B_{F^*}, e_1, \delta) = \emptyset$. Thus F star-denies the positive Daugavet property with $S(B_{F^*}, e_1, \delta) \cap S_{F^*}^+$. \square

We obtain the following fact by the successive application of Corollary 4.5 and Lemma 3.7.

Corollary 4.6. *Let $F \notin \mathfrak{N}_2$. Then $X_1 \oplus_F X_2$ is not a Daugavet range for any X_1 and X_2 .*

5. SUMS OF SPACES WHICH ARE NOT DAUGAVET DOMAINS

Lemma 5.1. *Let $F^* \notin \mathfrak{N}_2$. Then $X_1 \oplus_F X_2$ is not a Daugavet domain for any X_1 and X_2 .*

Proof. By Corollary 4.5 there is a $\delta > 0$ such that F^* star-denies the positive Daugavet property with $S(B_{F^{**}}, e_1, \delta) \cap S_{F^{**}}^+$. Recall that $F^{**} = F$. Therefore it follows from Lemma 3.5 that F denies the positive Daugavet property with $S(B_F, e_1, \delta) \cap S_F^+$. Then Lemma 3.6 gives the needed result. \square

We characterize the class of those F such that S_F^+ is a polygon with at most two edges, with the help of the following notation. Consider an F whose S_F^+ is a polygon with n edges. Denote $\hat{x}_1 := \max_{(1,y) \in S_F^+} y$ and $\hat{x}_2 := \max_{(x,1) \in S_F^+} x$. We say that F belongs to $\mathcal{F}_{n-1,n}$ if $\hat{x}_1 > 0$ and $\hat{x}_2 > 0$. If only one of \hat{x}_j equals zero, we say that $F \in \mathcal{F}_{n,n}$. And if both $\hat{x}_1 = \hat{x}_2 = 0$ then $F \in \mathcal{F}_{n+1,n}$ (see Figure 1).




FIGURE 1. Those F whose S_F^+ are presented on pictures (a) and (b), belong to $\mathcal{F}_{2,2}$. Picture (c) shows S_F^+ of $F \in \mathcal{F}_{3,2}$ and (d) shows S_F^+ of $F \in \mathcal{F}_{2,3}$.

Thus, $\mathfrak{N}_2 = \{\ell_1^{(2)}\} \cup \{\ell_\infty^{(2)}\} \cup \mathcal{F}_{2,2} \cup \mathcal{F}_{3,2}$. Let $n \in \mathbb{N}$ and $m \in \{n-1, n, n+1\}$. It is easy to see that $F^* \in \mathcal{F}_{n,m}$ if and only if $F \in \mathcal{F}_{m,n}$. Therefore, if $F^* \in \mathcal{F}_{2,2} \cup \mathcal{F}_{3,2}$ then $F \in \mathcal{F}_{2,2} \cup \mathcal{F}_{2,3}$. So, we obtain the following fact:

Corollary 5.2. *Let $F \notin \{\ell_1^{(2)}\} \cup \{\ell_\infty^{(2)}\} \cup \mathcal{F}_{2,2} \cup \mathcal{F}_{2,3} =: \mathfrak{M}_2$. Then $X_1 \oplus_F X_2$ is not a Daugavet domain for any X_1 and X_2 .*

6. EXAMPLES OF DAUGAVET CENTERS ACTING FROM AND INTO A SUM OF TWO BANACH SPACES

In this section we show that for every $F \in \mathfrak{M}_2$ there exists a Daugavet domain $X_1 \oplus_F X_2$, and for every $F \in \mathfrak{N}_2$ there is a Daugavet range $X_1 \oplus_F X_2$.

For $F = \ell_1^{(2)}$ and $F = \ell_\infty^{(2)}$ several examples of $X_1 \oplus_F X_2$ which are Daugavet domains and Daugavet ranges, are known. For instance, if X is a Daugavet domain then for every E the sum $X \oplus_\infty E$ is as well; and if X is a Daugavet range then $X \oplus_1 E$ is. If $G_1: X_1 \rightarrow Y_1$ and $G_2: X_2 \rightarrow Y_2$ are Daugavet centers then $G: X_1 \oplus_1 X_2 \rightarrow Y_1 \oplus_1 Y_2$ and $\tilde{G}: X_1 \oplus_\infty X_2 \rightarrow Y_1 \oplus_\infty Y_2$ which map every (x_1, x_2) into $(G_1 x_1, G_2 x_2)$, are Daugavet centers as well [3].

For future reference we mention the following fact:

Lemma 6.1 ([6], Lemma 2.8). *If X has the Daugavet property then for every finite-dimensional subspace Y_0 of X , every $\varepsilon > 0$, and every slice $S(B_X, x^*, \varepsilon)$ there is an $x \in S(B_X, x^*, \varepsilon)$ such that every $y \in Y_0$ and $t \in \mathbb{R}$ fulfill*

$$\|y + tx\| \geq (1 - \varepsilon)(\|y\| + |t|).$$

Consider an $F \in \mathcal{F}_{2,2} \cup \mathcal{F}_{2,3}$. Denote $c_1 := (1, \hat{x}_1) \in S_F^+$ and $c_2 := (\hat{x}_2, 1) \in S_F^+$. Then $[c_1, c_2] \subset S_F^+$ (see Figure 2).




FIGURE 2. Picture (a) shows S_F^+ of $F \in \mathcal{F}_{2,3}$ and (b) presents S_F^+ of $F \in \mathcal{F}_{2,2}$.

Let the line containing $[c_1, c_2]$ have the equation $f_1 a_1 + f_2 a_2 = 1$ with $(f_1, f_2) \in S_{F^*}^+$. Remark that for every $w^* \in S_{F^*}^+$ and $\varepsilon > 0$ we have $S(B_F, w^*, \varepsilon) \cap [c_1, c_2] \neq \emptyset$. In other words, there exists an $(a_1, a_2) \in S(B_F, w^*, \varepsilon)$ such that $f_1 a_1 + f_2 a_2 = 1$.

Proposition 6.2. *Let X have the Daugavet property, $F \in \mathcal{F}_{2,2} \cup \mathcal{F}_{2,3}$, and let $(f_1, f_2) \in S_{F^*}^+$ be the functional described above. Then $G: X \oplus_F X \rightarrow X$, $G(x_1, x_2) = f_1x_1 + f_2x_2$ is a Daugavet center.*

Proof. At first, calculate $\|G\|$:

$$\|G\| = \sup_{(x_1, x_2) \in S_{X \oplus_F X}} \|f_1x_1 + f_2x_2\| = \sup_{(x_1, x_2) \in S_{X \oplus_F X}} (f_1\|x_1\| + f_2\|x_2\|) = 1.$$

Let $\varepsilon > 0$, $y_0 \in S_X$ and $x^* = (x_1^*, x_2^*) \in S_{(X \oplus_F X)^*}$.

By Lemma 6.1 there exists an $\tilde{x}_1 \in B_X$ with $x_1^*(\tilde{x}_1) \geq \|x_1^*\|(1 - \varepsilon/4)$ and

$$(6.1) \quad \|y_0 + t\tilde{x}_1\| \geq \left(1 - \frac{\varepsilon}{4}\right) (1 + |t|)$$

for every $t \in \mathbb{R}$. Using again Lemma 6.1 we have an $\tilde{x}_2 \in B_X$ with $x_2^*(\tilde{x}_2) \geq \|x_2^*\|(1 - \varepsilon/4)$ and

$$(6.2) \quad \|y + t\tilde{x}_2\| \geq \left(1 - \frac{\varepsilon}{4}\right) (\|y\| + |t|)$$

for every $y \in \text{lin}\{y_0, \tilde{x}_1\}$ and every $t \in \mathbb{R}$.

Denote $w^* := (\|x_1^*\|, \|x_2^*\|) \in S_{F^*}^+$. Let $(a_1, a_2) \in S(B_F, w^*, 3\varepsilon/4)$ such that $f_1a_1 + f_2a_2 = 1$. Then for $x := (a_1\tilde{x}_1, a_2\tilde{x}_2) \in B_{X \oplus_F X}$ we have

$$\begin{aligned} x^*(x) &= a_1x_1^*(\tilde{x}_1) + a_2x_2^*(\tilde{x}_2) \geq \left(1 - \frac{\varepsilon}{4}\right) (a_1\|x_1^*\| + a_2\|x_2^*\|) \\ &\geq \left(1 - \frac{\varepsilon}{4}\right) \left(1 - \frac{3\varepsilon}{4}\right) > 1 - \varepsilon. \end{aligned}$$

Hence $x \in S(B_{X \oplus_F X}, x^*, \varepsilon)$ and

$$\|y_0 + Gx\| = \|y_0 + f_1a_1\tilde{x}_1 + f_2a_2\tilde{x}_2\|$$

by (6.2)

$$> \left(1 - \frac{\varepsilon}{4}\right) (\|y_0 + f_1a_1\tilde{x}_1\| + f_2a_2)$$

by (6.1)

$$> \left(1 - \frac{\varepsilon}{4}\right)^2 (1 + f_1a_1 + f_2a_2) = 2 \left(1 - \frac{\varepsilon}{4}\right)^2 > 2 - \varepsilon.$$

Theorem 1.2, item (ii) implies that G is a Daugavet center. \square

Corollary 6.3. *For an F there exists a Daugavet domain $X_1 \oplus_F X_2$ if and only if $F \in \mathfrak{M}_2$.*

Remark 6.4. *Note that $\mathfrak{M}_2 \not\subseteq \mathfrak{N}_2$. Then Corollary 6.3 and Corollary 4.6 imply that there exist Daugavet domains which are not Daugavet ranges.*

Now we present more examples of Daugavet centers acting from $X_1 \oplus_F X_2$ where $F = \ell_1^{(2)}$ or $F = \ell_\infty^{(2)}$.

Proposition 6.5. *Let X have the Daugavet property. Then*

- (a) *The operator $G: X \oplus_1 X \rightarrow X$, $G(x_1, x_2) = x_1 + x_2$ is a Daugavet center.*

- (b) For every $f_1, f_2 > 0$ the operator $G: X \oplus_\infty X \rightarrow X$, $G(x_1, x_2) = f_1 x_1 + f_2 x_2$ is a Daugavet center.

Proposition 6.5 can be proved the same way as Proposition 6.2.

Proposition 6.6. *Let X have the Daugavet property, $F \in \mathcal{F}_{2,2} \cup \mathcal{F}_{3,2}$, and let $(f_1, f_2) \in S_F^+$ be the vector described above. Then $G: X \rightarrow X \oplus_F X$, $Gx = (f_1 x, f_2 x)$ is a Daugavet center.*

Proof. Consider the adjoint operator $G^*: X^* \oplus_{F^*} X^* \rightarrow X^*$. For every $(x_1^*, x_2^*) \in X^* \oplus_{F^*} X^*$ and every $x \in X$ we have

$$G^*(x_1^*, x_2^*)(x) = \langle (f_1 x, f_2 x), (x_1^*, x_2^*) \rangle = f_1 x_1^*(x) + f_2 x_2^*(x).$$

Consequently, $G^*(x_1^*, x_2^*) = f_1 x_1^* + f_2 x_2^*$ for every $(x_1^*, x_2^*) \in X^* \oplus_{F^*} X^*$. By Proposition 6.2 G^* is a Daugavet center. The equation (1.1) implies that if G^* is a Daugavet center then G is as well. \square

Corollary 6.7. *For an F there exists a Daugavet range $X_1 \oplus_F X_2$ if and only if $F \in \mathfrak{N}_2$.*

Remark 6.8. *Since $\mathfrak{N}_2 \not\subseteq \mathfrak{M}_2$, we have the examples of Daugavet ranges which are not Daugavet domains.*

Proposition 6.9 which gives more examples of Daugavet centers acting into $X_1 \oplus_F X_2$ for $F = \ell_1^{(2)}$ and $F = \ell_\infty^{(2)}$, can be proved in a very similar way to Proposition 6.6.

Proposition 6.9. *Let X have the Daugavet property. Then*

- (a) *The operator $G: X \rightarrow X \oplus_\infty X$, $Gx = (x, x)$ is a Daugavet center.*
- (b) *For every $f_1, f_2 > 0$ the operator $G: X \rightarrow X \oplus_1 X$, $Gx = (f_1 x, f_2 x)$ is a Daugavet center.*

ACKNOWLEDGEMENTS

The author is grateful to her scientific adviser Vladimir M. Kadets for attention and numerous fruitful discussions, and to the referee for useful remarks.

REFERENCES

- [1] Bilik D., Kadets V., Shvidkoy R., Werner D., Narrow operators and the Daugavet property for ultraproducts, *Positivity*, 2005, 9, 45–62
- [2] Bourgain J., Rosenthal H.P., Martingales valued in certain subspaces of L^1 , *Israel J. Math.*, 1980, 37, 54–75
- [3] Bosenko T., Kadets V., Daugavet centers, *Zh. Mat. Fiz. Anal. Geom.*, (in press), preprint available at <http://arxiv.org/abs/0910.4503>
- [4] Daugavet I.K., On a property of completely continuous operators in the space C , *Uspekhi Mat. Nauk*, 1963, 18, 157–158, (in Russian)
- [5] Kadets V.M., Shepelska V., Werner D., Quotients of Banach spaces with the Daugavet property, *Bull. Pol. Acad. Sci. Math.*, 2008, 56, 131–147
- [6] Kadets V.M., Shvidkoy R.V., Sirotkin G.G., Werner D., Banach spaces with the Daugavet property, *Trans. Amer. Math. Soc.*, 2000, 352, 855–873

- [7] Kadets V.M., Werner D., A Banach space with the Schur and the Daugavet property, Proc. Amer. Math. Soc., 2004, 132, 1765–1773
- [8] Lindenstrauss J., Tzafriri L., Classical Banach Spaces II: Function spaces, Springer-Verlag, Berlin-Heidelberg-New York, 1979
- [9] Lozanovskii G.Ya., On almost integral operators in KB-spaces, Vestnik Leningrad Univ. Mat. Mekh. Astr., 1966, 21, 35–44, (in Russian)
- [10] Popov M.M., Daugavet type inequalities for narrow operators in the space L_1 , Mat. Stud., 2003, 20, 75–84
- [11] Shvidkoy R.V., Geometric aspects of the Daugavet property, J. Funct. Anal., 2000, 176, 198–212
- [12] Talagrand M., The three-space problem for L^1 , J. Amer. Math. Soc., 1990, 3, 9–29
- [13] Werner D., Recent progress on the Daugavet property, Irish Math. Soc. Bull., 2001, 46, 77–97
- [14] Werner D., The Daugavet equation for operators on function spaces, J. Funct. Anal., 1997, 143, 117–128
- [15] Wojtaszczyk P., Some remarks on the Daugavet equation, Proc. Amer. Math. Soc., 1992, 115, 1047–1052

DEPARTMENT OF MECHANICS AND MATHEMATICS, V.N. KARAZIN KHARKIV NATIONAL
UNIVERSITY, PL. SVOBODY 4, 61077 KHARKIV, UKRAINE
E-mail address: `t.bosenko@mail.ru`